

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, C07K 14/725, C12N 15/62, 5/10, 1/19		A3	(11) International Publication Number: WO 96/40918 (43) International Publication Date: 19 December 1996 (19.12.96)
(21) International Application Number: PCT/US96/09632 (22) International Filing Date: 6 June 1996 (06.06.96) (30) Priority Data: 08/477,733 7 June 1995 (07.06.95) US 08/484,624 7 June 1995 (07.06.95) US (71) Applicant: IMMUNEX CORPORATION [US/US]; 51 University Street, Seattle, WA 98101 (US). (72) Inventors: ARMITAGE, Richard, J.; 5133 Eagle Harbor Drive, Bainbridge Island, WA 98110 (US). FANSLOW, William, C., III; 218 S.W. 327th Place, Federal Way, WA 98023 (US). SPRIGGS, Melanie, K.; 2256 12th Avenue West, Seattle, WA 98119 (US). SRINIVASAN, Subhashini; 11325 N.E. 129th Street, Kirkland, WA 98034 (US). GIBSON, Marylou, G.; 7209 Wisteria Way, Carlsbad, CA 92009 (US). (74) Agent: PERKINS, Patricia, Anne; Immunex Corporation, 51 University Street, Seattle, WA 98101 (US).			(81) Designated States: AU, CA, IL, JP, KR, MX, NO, NZ, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 23 January 1997 (23.01.97)
(54) Title: CD40L MUTEIN			
(57) Abstract There is disclosed a polypeptide that is mutein of CD40L, and DNA sequences, vectors and transformed host cells useful in providing such CD40L muteins.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

NOVEL CD40L MUTEIN

BACKGROUND OF THE INVENTION

5 Human CD40 protein (CD40) is a peptide of 277 amino acids having a molecular weight of 30,600, and a 19 amino acid secretory signal peptide comprising predominantly hydrophobic amino acids (Stamenkovic et al., *EMBO J.* 8:1403, 1989). CD40 belongs to a family of receptors whose members include two forms of TNF receptor (Smith et al., *Science* 248:1019, 1990; and Schall et al., *Cell* 61:361, 1990), nerve growth factor receptor (Johnson et al., *Cell* 47:545, 1986), T cell antigen OX40 (Mallett et al., *EMBO J.* 9:1063, 1990), human *Fas* antigen (Itoh et al., *Cell* 66:233, 1991) and murine 4-1BB receptor (Kwon et al., *Cell. Immunol.* 121:414, 1989 [Kwon et al. I] and Kwon et al., *Proc. Natl. Acad. Sci. USA* 86:1963, 1989 [Kwon et al. II]). The molecular weight (exclusive of glycosylation) of the mature human CD40 protein is 28,300.

15 Activated CD4+ T cells express high levels of a ligand for CD40 (CD40L). Human CD40L, a membrane-bound glycoprotein, was cloned from peripheral blood T-cells as described in Spriggs et al., *J. Exp. Med.* 176:1543 (1992), and WO 93/08207. The cloning of murine CD40L is described in Armitage et al., *Nature* 357:80, 1992. CD40L induces B-cell proliferation in the absence of any co-stimulus, and can also induce production of immunoglobulins in the presence of cytokines. CD40L is a type II membrane polypeptide having an extracellular region at its C-terminus, a transmembrane region and an intracellular region at its N-terminus. Soluble forms of CD40L comprise the extracellular region of CD40L or a fragment thereof. CD40L biological activity is mediated by binding of the extracellular region of CD40L with CD40, and includes B cell proliferation and induction of antibody secretion (including IgE secretion).

25 Prior to the present invention, a ligand for CD40 that exhibited higher binding affinity for human CD40 than that of native CD40L was unknown. Accordingly, there is a need in the art to identify and characterize such a CD40 ligand mutein.

SUMMARY OF THE INVENTION

30 A novel cytokine, hereafter referred to as "CD40L," was isolated and characterized as described in WO 93/08207. The present invention comprises isolated DNA molecules encoding novel human CD40L muteins, and their complements. The isolated DNA molecules are selected from the group consisting of a DNA that encodes a polypeptide having an amino acid sequence as set forth in SEQ ID NO:2 wherein the cysteine at amino acid 194 is replaced with another amino acid, and DNA's that encode a polypeptide that is a fragment of the mutein. Preferably, the amino acid that is substituted for the Cys at residue

35

194 is selected from the group consisting of tryptophan, serine, aspartic acid, and lysine; most preferably, the amino acid is tryptophan.

The invention further provides DNA's encoding fragments of a CD40L mutein. The fragments are selected from the group consisting of peptides comprising amino acids 1 through 261, 35 through 261, 34 through 225, 113 through 261, 113 through 225, 120 through 261, or 120 through 225 of SEQ ID NO:2 in which the cysteine at the amino acid corresponding to amino acid 194 of SEQ ID NO:2 is replaced with another amino acid. DNA's that hybridize to any of the foregoing DNA's under stringent conditions (hybridization in 6 X SSC at 63°C overnight; washing in 3 X SSC at 55°C), and which encode a peptide that binds to CD40 and in which the cysteine at the amino acid corresponding to amino acid 194 of SEQ ID NO:2 is replaced with another amino acid, are also provided.

The invention further provides novel CD40L muteins encoded by the inventive DNA molecules. The novel muteins differ from native human CD40L in having an amino acid sequence as set forth in SEQ ID NO:2 wherein the cysteine at amino acid 194 is replaced with another amino acid. Preferably, the amino acid that is substituted for the Cys at residue 194 is selected from the group consisting of tryptophan, serine, aspartic acid, and lysine; most preferably, the amino acid is tryptophan. The novel muteins includes fragments of the extracellular domain of CD40L that bind CD40. Preferably the fragments are selected from the group consisting of peptides comprising amino acids 1 through 261, 35 through 261, 34 through 225, 113 through 261, 113 through 225, 120 through 261, or 120 through 225 of SEQ ID NO:2, most preferably comprising amino 113 through 261, in which the cysteine at the amino acid corresponding to amino acid 194 of SEQ ID NO:2 is replaced with another amino acid.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 presents the binding of trimeric human and murine CD40L and dimeric human and murine CD40L to C40/Fc as determined in a biosensor assay.

Figure 2 illustrates the binding of trimeric human CD40L (Figure 2A) and two preparations of monomeric human CD40L (Figure 2B) to C40/Fc in a biosensor assay.

DETAILED DESCRIPTION OF THE INVENTION

Novel polypeptides that can act as ligands for murine and human CD40 were isolated and sequenced as described in WO 93/08207. The present invention provides DNA molecules encoding novel human CD40L muteins, and their complements. The isolated DNA molecules are selected from the group consisting of a DNA that encodes a polypeptide having an amino acid sequence as set forth in SEQ ID NO:1 wherein the cysteine at amino acid 194 of SEQ ID NO:12 is substituted with an amino acid other than

cysteine, and DNA's that encode fragments thereof. Preferably, the amino acid that is substituted for the Cys are residue 194 is selected from the group consisting of tryptophan, serine, aspartic acid, and lysine; most preferably, the amino acid is tryptophan.

CD40L is a ligand for CD40, a receptor that is a member of the TNF receptor super family. Full-length CD40L is a membrane-bound polypeptide with an extracellular region at its C terminus, a transmembrane region, and an intracellular region at its N-terminus. A soluble version of CD40L can be made from the extracellular region or a fragment thereof. The extracellular region of human CD40L extends from amino acid 47 to amino acid 261 of SEQ ID NO:1. The biological activity of CD40L is mediated by binding to CD40 and comprises proliferation of B cells and induction of immunoglobulin secretion from activated B cells.

The novel CD40L mutein described herein was obtained by mutation of nucleotide sequences coding for a CD40L polypeptide. A CD40L mutein or analog, as referred to herein, is a polypeptide substantially homologous to a native CD40L but which has an amino acid sequence different from native sequence CD40L polypeptide because of one or a plurality of deletions, insertions or substitutions. Analogs of CD40L can be synthesized from DNA constructs prepared by oligonucleotide synthesis and ligation or by site-specific mutagenesis techniques.

Generally, substitutions should be made conservatively; i.e., the most preferred substitute amino acids are those which do not affect the ability of the inventive proteins to bind their receptors in a manner substantially equivalent to that of native CD40L. Examples of conservative substitutions include substitution of amino acids outside of the binding domain(s), and substitution of amino acids that do not alter the secondary and/or tertiary structure of CD40L. Additional examples include substituting one aliphatic residue for another, such as Ile, Val, Leu, or Ala for one another, or substitutions of one polar residue for another, such as between Lys and Arg; Glu and Asp; or Gln and Asn. Other such conservative substitutions, for example, substitutions of entire regions having similar hydrophobicity characteristics, are well known.

In another example, sequences encoding Cys residues can be altered to cause the Cys residues to be deleted or replaced with other amino acids, preventing formation of incorrect intramolecular disulfide bridges upon renaturation. Human CD40-L comprises five Cys residues in its extracellular domain. Thus, at least one of the five Cys residues can be replaced with another amino acid or deleted without effecting protein tertiary structure or disulfide bond formation. Preferably, the Cys that is substituted is amino acid 194 of SEQ ID NO:2, and the amino acid substituted therefor is selected from the group consisting of tryptophan, serine, aspartic acid, and lysine; most preferably, the amino acid is tryptophan.

Similarly, when a deletion or insertion strategy is adopted, the potential effect of the deletion or insertion on biological activity should be considered. Subunits of the inventive

proteins may be constructed by deleting terminal or internal residues or sequences to form fragments. Additional guidance as to the types of mutations that can be made is provided by a comparison of the sequence of CD40L to the sequences and structures of other TNF family members.

5 The primary amino acid structure of the inventive CD40L mutein may be modified to create CD40L derivatives by forming covalent or aggregative conjugates with other chemical moieties, such as glycosyl groups, lipids, phosphate, acetyl groups and the like, or by creating amino acid sequence mutants. Covalent derivatives of CD40L are prepared by linking particular functional groups to CD40L amino acid side chains or at the N-terminus or C-terminus of a CD40L mutein or the extracellular domain thereof.

10 Other derivatives of CD40L within the scope of this invention include covalent or aggregative conjugates of CD40L or its fragments with other proteins or polypeptides, such as by synthesis in recombinant culture as N-terminal or C-terminal fusions. For example, the conjugate may comprise a signal or leader polypeptide sequence at the N-terminal region or C-terminal region of a CD40L polypeptide which co-translationally or post-translationally directs transfer of the conjugate from its site of synthesis to a site inside or outside of the cell membrane or cell wall (e.g. the α -factor leader of *Saccharomyces*).

15 CD40L polypeptide fusions can comprise polypeptides added to facilitate purification and identification of CD40L (e.g. poly-His), or fusions with other cytokines to provide novel polyfunctional entities. Other cytokines include, for example, any of interleukins-1 through 13, TNF (tumor necrosis factor), GM-CSF (granulocyte macrophage-colony stimulating factor), G-CSF (granulocyte-colony stimulating factor), MGF (mast cell growth factor), EGF (epidermal growth factor), PDGF (platelet-derived growth factor), NGF (nerve growth factor), EPO (erythropoietin), γ -IFN (gamma interferon), 4-1BB-L (4-1BB ligand) and other cytokines that affect immune cell growth, differentiation or function.

20 Biological activity of CD40L may be determined, for example, by competition for binding to the ligand binding domain of CD40 (i.e. competitive binding assays). Several useful binding assays are disclosed herein. Those skilled in the art can readily apply routine experimentation to develop binding assays, using either isolated CD40 protein (i.e., CD40/Fc) or cells expressing CD40.

25 CD40L may also be assayed by measuring biological activity in a B cell proliferation assay. Human B cells may be obtained from human tonsils by purification by negative selection and Percoll density sedimentation, as described by Defrance et al., *J. Immunol.* 139:1135, 1987. Burkitt lymphoma cell lines may be used to measure cell proliferation in response to CD40L. Examples of Burkitt lymphoma cell lines include, for example, Raji (ATCC CCL 86), Daudi (ATCC CCL 213) and Namalwa (ATCC CRL 1432).

Yet another assay for determining CD40L biological activity is to measure immunoglobulin produced by B cells in response to activation by CD40L or a derivative or analog thereof. Polyclonal immunoglobulin secretion can be measured, for example, by incubating with 5×10^5 B cells/ml in culture for at least seven days. Immunoglobulin (Ig) production can be measured by an ELISA assay such as one described in Maliszewski et al., *J. Immunol.* 144:3028, 1990 [Maliszewski et al. I] or Maliszewski et al., *Eur J. Immunol.* 20:1735, 1990 [Maliszewski et al. II].

CD40L polypeptides may exist as oligomers, such as dimers or trimers. Oligomers are linked by disulfide bonds formed between cysteine residues on different CD40L polypeptides. Alternatively, one can link two soluble CD40L domains with a linker sequence such as those described in United States Patent 5,073,627, which is incorporated by reference herein. CD40L polypeptides may also be created by expression of a fusion protein comprising soluble CD40L (extracellular domain) and an Fc region of an immunoglobulin (for example, SEQ ID NO:4) to form divalent CD40L. Fusion proteins made with both heavy and light chains of an antibody will form a CD40L oligomer with as many as four CD40L extracellular regions. Trimeric CD40L is prepared by expressing a DNA encoding a fusion protein comprising a peptide that forms a trimer in solution (for example, a mutein of a yeast GCN4 "leucine zipper" (SEQ ID NO:5), or the lung surfactant protein D (SPD) trimerization domain (SEQ ID NO:6; Hoppe, et al., *FEBS Letters* 344:191, 1994) followed by the extracellular region of CD40L.

Fusion proteins can be prepared using conventional techniques of enzyme cutting and ligation of fragments from desired sequences. PCR techniques employing synthetic oligonucleotides may be used to prepare and/or amplify the desired fragments. Overlapping synthetic oligonucleotides representing the desired sequences can also be used to prepare DNA constructs encoding fusion proteins. Fusion proteins can also comprise CD40L and two or more additional sequences, including a leader (or signal peptide) sequence, oligomerizing region (i.e., Fc region, leucine zipper moiety or suitable zipper moiety), linker sequence, and sequences encoding highly antigenic moieties that provide a means for facile purification or rapid detection of a fusion protein.

Signal peptides facilitate secretion of proteins from cells. An exemplary signal peptide is amino acids -26 through -1 of SEQ ID NO:8. The Flag[®] octapeptide (Hopp et al., *Bio/Technology* 6:1204, 1988) does not alter the biological activity of fusion proteins, is highly antigenic and provides an epitope reversibly bound by a specific monoclonal antibody, enabling rapid detection and facile purification of the expressed fusion protein. The Flag[®] sequence is also specifically cleaved by bovine mucosal enterokinase at the residue immediately following the Asp-Lys pairing, fusion proteins capped with this peptide may also be resistant to intracellular degradation in *E. coli*. A murine monoclonal antibody that binds the Flag[®] sequence has been deposited with the ATCC under accession

number HB 9259; methods of using the antibody in purification of fusion proteins comprising the Flag[®] sequence are described in U.S. Patent 5,011,912, which is incorporated by reference herein.

Suitable Fc regions are defined as Fc regions that can bind to protein A or protein G, or alternatively, are recognized by an antibody that can be used in purification or detection of a fusion protein comprising the Fc region. Preferable Fc regions include the Fc region of human IgG₁ or murine IgG₁. One example is the human IgG₁ Fc region shown in SEQ ID NO:4. Fragments of a suitable Fc region may also be used, for example, an Fc region of human IgG₁ from which has been deleted a sequence of amino acids responsible for binding to protein A, such that the fragment binds to protein G but not protein A.

A linker sequence that separates the extracellular region of the CD40L from the oligomerizing moiety by a distance sufficient to ensure that the CD40L properly folds into its secondary and tertiary structures may also be used. Suitable linker sequences (1) will adopt a flexible extended conformation, (2) will not exhibit a propensity for developing an ordered secondary structure which could interact with the functional domains of fusion proteins, and (3) will have minimal hydrophobic or charged character which could promote interaction with the functional protein domains. Typical surface amino acids in flexible protein regions include Gly, Asn and Ser. Virtually any permutation of amino acid sequences containing Gly, Asn and Ser would be expected to satisfy the above criteria for a linker sequence. Other near neutral amino acids, such as Thr and Ala, may also be used in the linker sequence. The length of the linker sequence may vary without significantly affecting the biological activity of the fusion protein. Linker sequences are unnecessary where the proteins being fused have non-essential N- or C-terminal amino acid regions which can be used to separate the functional domains and prevent steric interference.

DNA constructs that encode various additions or substitutions of amino acid residues or sequences, or deletions of terminal or internal residues or sequences not needed for biological activity or binding can be prepared. For example, the extracellular CD40L N-glycosylation site can be modified to preclude glycosylation while allowing expression of a homogeneous, reduced carbohydrate analog using yeast expression systems. N-glycosylation sites in eukaryotic polypeptides are characterized by an amino acid triplet Asn-X-Y, wherein X is any amino acid except Pro and Y is Ser or Thr. Appropriate modifications to the nucleotide sequence encoding this triplet will result in substitutions, additions or deletions that prevent attachment of carbohydrate residues at the Asn side chain.

Other approaches to mutagenesis involve modification of sequences encoding dibasic amino acid residues to enhance expression in yeast systems in which KEX2 protease activity is present. Sub-units of a CD40L polypeptide may be constructed by deleting sequences encoding terminal or internal residues or sequences. Moreover, other

analyses may be performed to assist the skilled artisan in selecting sites for mutagenesis. For example, WO 93/08207 (the disclosure of which is hereby incorporated by reference) discusses methods of selecting ligand agonists and antagonists.

5 The sequence of murine CD40L cDNA was obtained by direct expression techniques; the sequence of human CD40L was obtained by cross-species hybridization techniques using the murine CD40L cDNA as a probe. Both are described in WO 93/08207. Briefly, the extracellular region of human CD40 (the receptor; SEQ ID NO:3) was cloned by polymerase chain reaction (PCR) techniques using primers based upon a sequence published in Stamenkovic et al. A CD40/Fc fusion protein was obtained by PCR
10 techniques by fusing DNA encoding the extracellular domain of CD40 with DNA encoding the Fc domain of human IgG1 (SEQ ID NO:4). Purified soluble CD40 proteins were able to antagonize CD40L-mediated biological activities.

A cDNA library was prepared from an EL4 cell line sorted by FACS (fluorescence activated cell sorting) on the basis of binding of a biotinylated CD40/Fc fusion protein.
15 Cells were sorted five times until there was a significant shift in fluorescence intensity based upon expression of a ligand for CD40 by the sorted EL-4 cells. Briefly, cDNA was synthesized, inserted into empty pDC406 vector and transformed into *E. coli*. Transformants were pooled, and the DNA from the pools was isolated and transfected into CV1-EBNA cells to create an expression cloning library. Transfected CV1-EBNA cells
20 were cultured on slides to permit transient expression of CD40L. The slides were screened with radioiodinated CD40/Fc, and examined to identify cells expressing CD40L by ascertaining the presence of autoradiographic silver grains against a light background.

A single clone was isolated and sequenced by standard techniques, to provide the cDNA sequence and deduced amino acid sequence of murine CD40L. The human
25 homologue CD40L cDNA was found by cross species hybridization techniques. Briefly, a human peripheral blood lymphocyte (PBL) cDNA library was made from activated peripheral blood lymphocytes. A murine probe was constructed corresponding to the coding region of murine CD40L from nucleotide 13 to nucleotide 793 of murine CD40L. Using standard techniques for cross-species hybridization, the probe was used to identify a
30 clone that expressed human CD40L. The nucleotide and amino acid sequences of human CD40L are shown in SEQ ID NO's 1 and 2.

Recombinant expression vectors for expression of CD40L by recombinant DNA techniques include a CD40L DNA sequence comprising a synthetic or cDNA-derived DNA fragment encoding a CD40L polypeptide, operably linked to a suitable transcriptional or
35 translational regulatory nucleotide sequence, such as one derived from a mammalian, microbial, viral, or insect gene. Examples of regulatory sequences include sequences having a regulatory role in gene expression (e.g., a transcriptional promoter or enhancer), optionally an operator sequence to control transcription, a sequence encoding an mRNA

ribosomal binding site, and appropriate sequences which control transcription and translation initiation and termination.

Nucleotide sequences are operably linked when the regulatory sequence functionally relates to the CD40L DNA sequence. Thus, a promoter nucleotide sequence is operably
5 linked to a CD40L DNA sequence if the promoter nucleotide sequence controls the transcription of the CD40L DNA sequence. Still further, a ribosome binding site may be operably linked to a sequence for a CD40L polypeptide if the ribosome binding site is positioned within the vector to encourage translation. In addition, sequences encoding signal peptides can be incorporated into expression vectors. For example, a DNA sequence
10 for a signal peptide (secretory leader) may be operably linked to a CD40L DNA sequence

Suitable host cells for expression of CD40L polypeptides include prokaryotes, yeast or higher eukaryotic cells. Prokaryotes include gram negative or gram positive organisms, for example, *E. coli* or *Bacilli*. Suitable prokaryotic host cells for transformation include, for example, *E. coli*, *Bacillus subtilis*, *Salmonella typhimurium*, and various other species
15 within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*. Higher eukaryotic cells include established cell lines of mammalian origin. Cell-free translation systems could also be employed to produce CD40L polypeptides using RNAs derived from DNA constructs disclosed herein. Appropriate cloning and expression vectors for use with bacterial, fungal, yeast, and mammalian cellular hosts are described, for example, in
20 Pouwels et al. *Cloning Vectors: A Laboratory Manual*, Elsevier, New York, (1985).

The expression vectors carrying the recombinant CD40L DNA sequence are transfected or transformed into a substantially homogeneous culture of a suitable host microorganism or mammalian cell line. Transformed host cells are cells which have been transformed or transfected with nucleotide sequences encoding CD40L polypeptides and
25 express CD40L polypeptides. Expressed CD40L polypeptides will be located within the host cell and/or secreted into culture supernatant fluid, depending upon the nature of the host cell and the gene construct inserted into the host cell.

Expression vectors transfected into prokaryotic host cells generally comprise one or more phenotypic selectable markers. A phenotypic selectable marker is, for example, a
30 gene encoding a protein that confers antibiotic resistance or that supplies an autotrophic requirement, and an origin of replication recognized by the host to ensure amplification within the host. Other useful expression vectors for prokaryotic host cells include a selectable marker of bacterial origin derived from commercially available plasmids. This selectable marker can comprise genetic elements of the cloning vector pBR322 (ATCC
35 37017). pBR322 contains genes for ampicillin and tetracycline resistance and thus provides simple means for identifying transformed cells. The pBR322 "backbone" sections are combined with an appropriate promoter and a CD40L DNA sequence. Other

commercially available vectors include, for example, pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden) and pGEM1 (Promega Biotec, Madison, WI, USA).

Promoter sequences are commonly used for recombinant prokaryotic host cell expression vectors. Common promoter sequences include β -lactamase (penicillinase),
5 lactose promoter system (Chang et al., *Nature* 275:615, 1978; and Goeddel et al., *Nature* 281:544, 1979), tryptophan (trp) promoter system (Goeddel et al., *Nucl. Acids Res.* 8:4057, 1980; and EP-A-36776) and tac promoter (Maniatis, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, p. 412, 1982). A particularly useful
10 prokaryotic host cell expression system employs a phage λ P_L promoter and a cI857ts thermolabile repressor sequence. Plasmid vectors available from the American Type Culture Collection which incorporate derivatives of the λ P_L promoter include plasmid pHUB2 (resident in *E. coli* strain JMB9 (ATCC 37092)) and pPLc28 (resident in *E. coli* RR1 (ATCC 53082)).

CD40L may be expressed in yeast host cells, preferably from the *Saccharomyces*
15 genus (e.g., *S. cerevisiae*). Other genera of yeast, such as *Pichia* or *Kluyveromyces*, may also be employed. Yeast vectors will often contain an origin of replication sequence from a 2 μ yeast plasmid, an autonomously replicating sequence (ARS), a promoter region, sequences for polyadenylation, and sequences for transcription termination. Preferably, yeast vectors include an origin of replication sequence and selectable marker. Suitable
20 promoter sequences for yeast vectors include promoters for metallothionein, 3-phosphoglycerate kinase (Hitzeman et al., *J. Biol. Chem.* 255:2073, 1980) or other glycolytic enzymes (Hess et al., *J. Adv. Enzyme Reg.* 7:149, 1968; and Holland et al., *Biochem.* 17:4900, 1978), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate
25 isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase. Other suitable vectors and promoters for use in yeast expression are further described in Hitzeman, EPA-73,657.

Mammalian or insect host cell culture systems could also be employed to express recombinant CD40L polypeptides. Examples of suitable mammalian host cell lines include
30 the COS-7 line of monkey kidney cells (ATCC CRL 1651) (Gluzman et al., *Cell* 23:175, 1981), L cells, C127 cells, 3T3 cells (ATCC CCL 163), Chinese hamster ovary (CHO) cells, HeLa cells, and BHK (ATCC CRL 10) cell lines. Suitable mammalian expression vectors include nontranscribed elements such as an origin of replication, a promoter sequence, an enhancer linked to the structural gene, other 5' or 3' flanking nontranscribed
35 sequences, such as ribosome binding sites, a polyadenylation site, splice donor and acceptor sites, and transcriptional termination sequences.

Transcriptional and translational control sequences for mammalian host cell expression vectors may be excised from viral genomes. For example, commonly used

mammalian cell promoter sequences and enhancer sequences are derived from Polyoma virus, Adenovirus 2, Simian Virus 40 (SV40), and human cytomegalovirus. DNA sequences derived from the SV40 viral genome, for example, SV40 origin, early and late promoter, enhancer, splice, and polyadenylation sites may be used to provide the other genetic elements required for expression of a structural gene sequence in a mammalian host cell. Viral early and late promoters are particularly useful because both are easily obtained from a viral genome as a fragment which may also contain a viral origin of replication (Fiers et al., *Nature* 273:113, 1978). Smaller or larger SV40 fragments may also be used, provided the approximately 250 bp sequence extending from the *Hind* III site toward the *Bgl* I site located in the SV40 viral origin of replication site is included.

Exemplary mammalian expression vectors can be constructed as disclosed by Okayama and Berg (*Mol. Cell. Biol.* 3:280, 1983). A useful high expression vector, PMLSV N1/N4, described by Cosman et al., *Nature* 312:768, 1984 has been deposited as ATCC 39890. Additional useful mammalian expression vectors are described in EP-A-0367566, and in U.S. Patent Application Serial No. 07/701,415, filed May 16, 1991, incorporated by reference herein. For expression of a type II protein extracellular region, such as CD40L, a heterologous signal sequence should be added, such as the signal sequence for interleukin-7 (IL-7) described in United States Patent 4,965,195, or the signal sequence for interleukin-2 receptor described in United States Patent Application 06/626,667 filed on July 2, 1984.

Purification of Recombinant CD40L Polypeptides

CD40L polypeptides may be prepared by culturing transformed host cells under culture conditions necessary to express CD40L polypeptides. The resulting expressed polypeptides may then be purified from culture media or cell extracts. A CD40L polypeptide, if desired, may be concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. Following the concentration step, the concentrate can be applied to a purification matrix such as a gel filtration medium. Alternatively, an anion exchange resin can be employed, for example, a matrix or substrate having pendant diethylaminoethyl (DEAE) groups. The matrices can be acrylamide, agarose, dextran, cellulose or other types commonly employed in protein purification. Alternatively, a cation exchange step can be employed. Suitable cation exchangers include various insoluble matrices comprising sulfopropyl or carboxymethyl groups. Sulfopropyl groups are preferred.

Finally, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, (e.g., silica gel having pendant methyl or other aliphatic groups) can be employed to further purify CD40L. Some or all of

the foregoing purification steps, in various combinations, can also be employed to provide a substantially homogeneous recombinant protein.

It is also possible to utilize an affinity column comprising CD40 ligand binding domain to affinity-purify expressed CD40L polypeptides. CD40L polypeptides can be removed from an affinity column in a high salt elution buffer and then dialyzed into a lower salt buffer for use.

Recombinant protein produced in bacterial culture is usually isolated by initial disruption of the host cells, centrifugation, extraction from cell pellets if an insoluble polypeptide, or from the supernatant fluid if a soluble polypeptide, followed by one or more concentration, salting-out, ion exchange, affinity purification or size exclusion chromatography steps. Finally, RP-HPLC can be employed for final purification steps. Microbial cells can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents.

Transformed yeast host cells are preferably employed to express CD40L as a secreted polypeptide. This simplifies purification. Secreted recombinant polypeptide from a yeast host cell fermentation can be purified by methods analogous to those disclosed by Urdal et al. (*J. Chromatog.* 296:171, 1984). Urdal et al. describe two sequential, reversed-phase HPLC steps for purification of recombinant human IL-2 on a preparative HPLC column.

Administration of CD40L Compositions

The present invention provides therapeutic compositions comprising an effective amount of CD40L in a suitable diluent or carrier and methods of treating mammals using the compositions. For therapeutic use, purified CD40L or a biologically active analog thereof is administered to a patient, preferably a human, for treatment in a manner appropriate to the indication. Thus, for example, CD40L pharmaceutical compositions (for example, in the form of a soluble extracellular domain, or a fragment thereof) which is administered to achieve a desired therapeutic effect can be given by bolus injection, continuous infusion, sustained release from implants, or other suitable technique.

Typically, a CD40L therapeutic agent will be administered in the form of a pharmaceutical composition comprising purified CD40L polypeptide in conjunction with physiologically acceptable carriers, excipients or diluents. Such carriers will be nontoxic to patients at the dosages and concentrations employed. Ordinarily, the preparation of such compositions entails combining a CD40L polypeptide with buffers, antioxidants such as ascorbic acid, low molecular weight (less than about 10 residues) polypeptides, proteins, amino acids, carbohydrates including glucose, sucrose or dextrans, chelating agents such as EDTA, glutathione and other stabilizers and excipients. Neutral buffered saline or saline mixed with conspecific serum albumin are exemplary appropriate diluents.

The following examples are intended to illustrate particular embodiments and not limit the scope of the invention. The relevant disclosures of all publications cited herein are specifically incorporated by reference.

EXAMPLE 1

This example describes two solid-phase binding assays, the first of which, (a), can be used to assess the ability of trimeric CD40L to bind CD40, and the second of which, (b), is used to detect the presence of CD40L.

(a) *Quantitative CD40L ELISA*

CD40/Fc is prepared and purified as described in WO 93/08207, and used to coat 96-well plates (Corning EasyWash ELISA plates, Corning, NY, USA). The plates are coated with 2.5 µg/well of CD40/Fc in PBS overnight at 4°C, and blocked with 1% non-fat milk in PBS for 1 hour at room temperature. Samples to be tested are diluted in 10% normal goat serum in PBS, and 50 µl is added per well. A titration of unknown samples is run in duplicate, and a titration of reference standard of CD40L is run to generate a standard curve. The plates are incubated with the samples and controls for 45 minutes at room temperature, then washed four times with PBS. Second step reagent, rabbit anti-oligomerizing zipper, is added (50 µl/well, concentration approximately 2.5 µg/ml), and the plates are incubated at room temperature for 45 minutes. The plates are again washed as previously described, and goat F(ab')₂ anti-rabbit IgG conjugated to horseradish peroxidase (Tago, Burlingame, CA, USA) is added. Plates are incubated for 45 minutes at room temperature, washed as described, and the presence of CD40L is detected by the addition of chromogen, tetramethyl benzidine (TMB; 100µl/well) for 15 minutes at room temperature. The chromogenic reaction is stopped by the addition of 100 µl/well 2N H₂SO₄, and the OD₄₅₀-OD₅₆₂ of the wells determined. The quantity of trimeric CD40L can be determined by comparing the OD values obtained with the unknown samples to the values generated for the standard curve. Values are expressed as the number of binding units per ml. A binding unit is roughly one ng of protein as estimated using a purified Fc fusion protein of the ligand as a standard. In this manner, the concentration and specific activity of several different batches of trimeric CD40L have been determined.

(b) *Qualitative Dot Blot*

CD40L trimer (1 µl of crude supernatant or column fractions) is adsorbed to dry BA85/21 nitrocellulose membranes (Schleicher and Schuell, Keene, NH) and allowed to dry. The membranes are incubated in tissue culture dishes for one hour in Tris (0.05 M) buffered saline (0.15 M) pH 7.5 containing 1% w/v BSA to block nonspecific binding sites. At the end of this time, the membranes are washed three times in PBS, and rabbit anti-oligomerizing zipper antibody is added at an approximate concentration of 10 µg/ml in PBS containing 1% BSA, following which the membranes are incubated for one hour at

room temperature. The membranes are again washed as described, and a horseradish peroxidase (HRP)-labeled antibody (such as goat anti-rabbit Ig; Southern Biotech, Birmingham, AL) at an approximate dilution of 1:1000 in PBS containing 1% BSA is added. After incubating for one hour at room temperature, the membranes are washed and
5 chromogen (i.e. 4-chloronaphthol reagent, Kirkegard and Perry, Gaithersburg, MD) is added. Color is allowed to develop for ten minutes at room temperature, and the reaction is stopped by rinsing the membranes with water. The membranes are washed, and the presence of CD40L is determined by analyzing for the presence of a blue-black color. This assay was used to determine the presence or absence of trimeric CD40L in cell culture
10 supernatant fluids and in purification column fractions. The assay further provides a semi-quantitative method of determining relative amounts of trimeric CD40L by comparing the intensity of the color in unknown samples to the intensity of known quantities of controls.

EXAMPLE 2

15 This example describes construction of a human CD40L DNA construct to express trimeric CD40L in Chinese hamster ovary (CHO) cells. Trimeric CD40L contains a leader sequence, and a 33 amino acid sequence referred to as an oligomerizing zipper (SEQ ID NO:5), followed by the extracellular region of human CD40L from amino acid 51 to amino acid 261 (SEQ ID NO:1). The construct was prepared by cutting the appropriate DNA
20 from a plasmid containing human CD40L, and ligating the DNA into the expression vector pCAVDHFR. The resultant construct was referred to as CAV/DHFR-CD40LT. pCAVDHFR includes regulatory sequences derived from cytomegalovirus, SV40, and Adenovirus 2, along with the gene for dihydrofolate reductase (DHFR), and allows random integration of a desired gene into host cell chromosomes. Expression of DHFR enables
25 the DHFR⁻ host cells to grow in media lacking glycine, hypoxanthine, and thymidine (GHT). A similar construct was also made for expression of murine CD40L timer in CHO cells. In addition to the leader and oligomerizing zipper sequences, the murine construct also contained a sequence encoding the octapeptide referred to as Flag® (described previously) between the trimerization domain ("leucine zipper" or oligomerizing zipper) and
30 the extracellular region of murine CD40L. The nucleotide and amino acid sequence of the human CD40L-encoding DNA's are shown in SEQ ID NO's 7 and 8 respectively. Additional constructs can be prepared using standard methods. For example, vectors which incorporates dual promoters such as those described in U.S. Patent 4,656,134, or vectors employing enhancer sequences such as those described in U.S. Patent 4,937,190
35 or in Kaufman et al., *Nucl. Acids Res.* 19:4485, 1991, are also useful in preparing constructs for expression of CD40L in CHO cells.

The resulting ligation product was transfected into CHO cells using either Lipofectin® Reagent or Lipofectamine™ Reagent (Gibco BRL, Gaithersburg, MD). Both

of these reagents are commercially available reagents used to form lipid-nucleic acid complexes (or liposomes) which, when applied to cultured cells, facilitate uptake of the nucleic acid into the cells. Cells which were transfected with the pCAVDHFR-CD40LT construct were selected in DMEM:F12 medium in the absence of GHT. Cells which were
5 able to grow in the absence of GHT were tested for production of CD40L using a solid phase binding assay as described in Example 16. Results indicated that in this transfection system, Lipofectamine™ Reagent gave higher rates of successful transfection.

Approximately 160 clones were screened and two positive clones were identified and expanded for further study. Cells were passaged in GHT-free DMEM:F12 medium,
10 and monitored for stability by assessing production of trimeric CD40L in the solid-phase binding assay described above. Based on these results, one clone was chosen which appeared to be stably transfected with the CD40L DNA, and which produced and secreted approximately $1 \mu\text{g}/10^6$ cells/day of CD40L trimer. Additional constructs comprising other vectors and all or a portion of the DNA sequences described in this example can be
15 used to prepare additional stably transfected cell lines, substantially as described herein.

Once such stably transfected cells were identified, large scale cultures of transfected cells were grown to accumulate supernatant containing trimeric CD40L. Suitable large-scale culture conditions include the use of bioreactors. Similar procedures were followed to produce CHO cell lines that secreted a trimeric murine CD40L at
20 approximately $0.05 \mu\text{g}/10^6$ cells/day. CHO cells stably transfected with either the human or murine CD40L construct, having acquired a DHFR gene from the pCAVDHFR plasmid, are resistant to methotrexate. Methotrexate can be added to the culture medium to amplify the number of copies of the CD40L trimer DNA in order to increase production of CD40L
25 trimer.

EXAMPLE 3

This example illustrates the binding affinities of several different CD40L constructs. Affinity experiments were conducted by biospecific interaction analysis (BIA) using a biosensor, an instrument that combines a biological recognition mechanism with a sensing
30 device or transducer. An exemplary biosensor is BIAcore™, from Pharmacia Biosensor AB (Uppsala, Sweden; see Fägerstam L.G., *Techniques in Protein Chemistry II*, ed. J.J. Villafranca, Acad. Press, NY, 1991). BIAcore™ uses the optical phenomenon surface plasmon resonance (Kretschmann and Raether, *Z. Naturforschung, Teil. A* 23:2135, 1968) to monitor the interaction of two biological molecules. Molecule pairs having affinity
35 constants in the range 10^5 to 10^{10} M^{-1} , and association rate constants in the range of 10^3 to $10^6 \text{ M}^{-1}\text{s}^{-1}$, are suitable for characterization with BIAcore™.

The biosensor chips were coated with goat anti-human IgG₁ Fc, which was used to bind CD40/Fc to the chip. The different constructs of CD40L were then added at

increasing concentrations; the chip was regenerated between the different constructs by the addition of sodium hydroxide. Two separate experiments were performed. In the first, the binding of a dimeric human CD40L (CD40L/Fc), trimeric human CD40L (CD40L/"leucine zipper"), and dimeric and trimeric murine CD40L (prepared substantially as described for the human CD40L) were compared. In the second experiment, the binding of trimeric human CD40L was compared to the binding of two different preparations of monomeric human CD40L (comprising solely the portion of the extracellular domain most homologous to TNF). The resultant data were analyzed to determine the affinity and association rate constants of the different CD40L constructs. Results are shown in Table 1 below, and in Figures 1 and 2.

Table 1: Binding of CD40L to CD40/Fc

	Site 1 (mol/mol CD40/Fc)	K ₁ (M ⁻¹)	Site2 (mol/mol CD40/Fc)	K ₂ (M ⁻¹)
Human Trimer	0.04 ± 0.02	5 ± 3 x 10 ⁹	0.68 ± 0.07	7.5 ± 2.5 x 10 ⁷
Human Dimer	0.013 ± 0.002	1.6 ± 0.7 x 10 ¹¹	0.049 ± 0.004	7.0 ± 2.1 x 10 ⁸
Murine Trimer	0.02 ± 0.003	6 ± 3 x 10 ¹⁰	0.44 ± 0.02	1.6 ± 0.2 x 10 ⁸
Murine Dimer	0.05 ± 0.02	7 ± 4 x 10 ⁹	0.14 ± 0.23	4.0 ± 1.0 x 10 ⁷
Human Trimer	0.26	6.6 x 10 ⁸	0.41	2.6 x 10 ⁷
Monomer #1	Not Detected	Not Detected	1.20	4.6 x 10 ⁷
Monomer #2	Not Detected	Not Detected	1.27	1.1 x 10 ⁷

Analysis of the data indicated that a CD40L monomer comprising solely the portion of the extracellular domain most homologous to TNF was capable of binding CD40, although with somewhat lower affinity than oligomeric CD40L. An analysis of the ratio of binding in the second experiment demonstrated that there are twice as many CD40L monomer units bound per CD40/Fc molecule as trimeric CD40L, confirming that two monomers of CD40L bind one CD40/Fc dimer and one trimeric CD40L binds one CD40/Fc dimer.

Example 4

This example illustrates preparation of a number of muteins of a CD40 ligand/zipper domain fusion protein. Mutations for constructs to be expressed in yeast (mutants 14, 18, 32, 41, 43, 10PP and 18PP) were generated by PCR misincorporation (Mulrad et al *Yeast* 8:79, 1992), and selected based on an apparent increase in secretion as improved secretion mutants. Mutants 14, 18, 32, 41, and 43 were isolated in *S. cerevisiae*. Mutants 10PP and 18PP were isolated in *P. pastoris*. Mutations for constructs to be expressed in mammalian cells (FL194.W, 194.W, LZ12V, 215.T, 255.F, and 194.S) were also prepared using PCR, and were either the result of site-directed mutagenesis or were the random product of

PCR. The types of mutations obtained and their effect on activity (ability to bind CD40 in a solid phase binding assay substantially as described in Example 1) are shown in Table 2 below.

Table 2: Mutations present in the CD40 ligand/zipper domain fusion protein

Mutant No.:	Zipper Domain Mutation ^a	CD40L Domain Mutations ^b	Activity	Type of Mutant
14	I12N	K260N	+	random mutant
18	L13P	A130P, R181Q	+	random mutant
32	I12N	Q121P	+	random mutant
41	I5M, I16T	NA	+	random mutant
43	I16N	T134S, K164I, Q186L, N210S	+	random mutant
10PP	I9N, K27R	NA ^d	+	random mutant
18PP	L13P	NA	+	random mutant
LZ12.V	I12V	Deletion of aa 1-112	+	PCR; random
215.T	NA	Deletion of aa 1-112; A215T	+	PCR; random
255.F	NA	Deletion of aa 1-112; S255F	-	PCR; site-directed
FL194W	NA	C194W	+	PCR; site-directed
194.W	NA	Deletion of aa 1-112; C194W	+	PCR; site-directed
194.S	NA	Deletion of aa 1-112; C194S	NDe	PCR; site-directed
194.A	NA	Deletion of aa 1-112; C194A	NDe	PCR; site-directed
194.D	NA	Deletion of aa 1-112; C194D	NDe	PCR; site-directed
194.K	NA	Deletion of aa 1-112; C194K	NDe	PCR; site-directed

5 a: Mutations are given as the residue present in the native peptide, the residue number, and the residue present in the mutein. Residue numbers for zipper domain mutations are relative to SEQ ID NO:5.

b: Residue numbers for mutations in the CD40L domain are relative to SEQ ID NO:1.

c: Mutant 10PP also contained mutations in regions other than CD40L domain or the zipper domain (T-4S, D-2P, relative to SEQ ID NO:7).

10 d: Not applicable

e: Not done

Mutant 18PP had only a single mutation in the molecule, which was sufficient to affect secretion in yeast. Mutant 41 had two mutations, both of which were in the isoleucine residues of the zipper domain. The mutations in the zipper improve secretion from yeast without apparent effect on activity. Mutant 194.W was expressed in yeast cells and purified either by a combination of ion exchange chromatography steps (194.W (c)) or by affinity chromatography (194.W (a)) using a monoclonal antibody that binds the oligomerizing zipper moiety. oligomerizing zipper moiety. The yeast-expressed mutant (194.W) exhibited greater affinity for CD40 in a biosensor assay performed substantially as described in Example 3, and exhibited greater biological

activity than wild type CD40 ligand/zipper domain fusion protein (WT) expressed in yeast, in a B cell proliferation assay. These results are shown in Table 3.

Table 3: Comparison of WT and 194.W for Receptor Binding and B-cell Proliferation

	Affinity K_a (M^{-1})	B-cell proliferation ($U/\mu g^a$)	
		Experiment 1	Experiment 2 ^d
WT	7.7×10^7	77 ^b	15
194.W (c)	1.8×10^9	171	116
194.W (a)	ND ^c	ND	161

a: A unit (U) is the concentration that induces half-maximal proliferation

b: Average from two independent preparations

c: Not done

d: Average of two assays

Moreover, FL194W expressed in mammalian cells also demonstrated higher binding than WT CD40L in a semi-quantitative western blot analysis.

Additional constructs were prepared by substituting the lung surfactant protein D (SPD) trimerization domain (SEQ ID NO:6; Hoppe, et al., *FEBS Letters* 344:191, 1994) in place of the trimer-forming zipper of SEQ ID NO:5. This construct is expressed in *S. cerevisiae* and in mammalian cells at low levels. Activity is determined as described previously; various mutants based on such constructs can also be prepared to optimize secretion or other product characteristics, as described above.

EXAMPLE 5

This example illustrates the binding affinities of several different CD40L constructs. Affinity experiments were conducted by biospecific interaction analysis (BIA), substantially as described in Example 3, using two of the constructs described in Example 4. Results presented in Table 4 represent the average values obtained for two different preparations of wild-type (WT) CD40LT, and three different preparations of mutein (194.W). Results are shown below.

Table 4: Binding of CD40LT to CD40/Fc

	K_1 (M^{-1})	K_2 (M^{-1})
CD40LT WT	$7.4 \pm 3.6 \times 10^8$	$5.4 \pm 1.5 \times 10^8$
CD40LT 194.W	$9.7 \pm 3.6 \times 10^9$	$3.0 \pm 0.4 \times 10^7$

These results confirm that the CD40LT 194.W has a higher affinity for CD40/Fc than does the wild type CD40LT.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- 5 (i) APPLICANT: IMMUNEX CORPORATION
(ii) TITLE OF INVENTION: NOVEL CD40L MUTEIN
(iii) NUMBER OF SEQUENCES: 8
(iv) CORRESPONDENCE ADDRESS:
10 (A) ADDRESSEE: IMMUNEX CORPORATION
(B) STREET: 51 UNIVERSITY STREET
(C) CITY: SEATTLE
(D) STATE: WASHINGTON
(E) COUNTRY: USA
(F) ZIP: 98101
15
(v) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: Apple Macintosh
(C) OPERATING SYSTEM: Apple Operating System 7.5.3
20 (D) SOFTWARE: Microsoft Word for Apple, version 6.0.1a

(vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER:
(B) FILING DATE: : 06 JUN 1996
25 (C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: USSN 08/477,733
(B) FILING DATE: 07 JUN 1995
30 (C) CLASSIFICATION:

(vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: USSN 08/484,624
(B) FILING DATE: 07 JUN 1995
35 (C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:
(A) NAME: Perkins, Patricia Anne
40 (B) REGISTRATION NUMBER: 34,693
(C) REFERENCE/DOCKET NUMBER: 2802-DWO

(ix) TELECOMMUNICATION INFORMATION:
(A) TELEPHONE: 2065870430
45 (B) TELEFAX: 2062330644

(2) INFORMATION FOR SEQ ID NO:1:

- 50 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 840 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
55 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA
(iii) HYPOTHETICAL: NO
(iv) ANTI-SENSE: NO

60 (vi) ORIGINAL SOURCE:
(A) ORGANISM: Homo sapiens

(vii) IMMEDIATE SOURCE:
(B) CLONE: CD40L

5 (ix) FEATURE:
(A) NAME/KEY: CDS
(B) LOCATION: 46..831

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

10 TGCCACCTTC TCTGCCAGAA GATACCATTT CAACTTTAAC ACAGC ATG ATC GAA 54
Met Ile Glu
1

15 ACA TAC AAC CAA ACT TCT CCC CGA TCT GCG GCC ACT GGA CTG CCC ATC 102
Thr Tyr Asn Gln Thr Ser Pro Arg Ser Ala Ala Thr Gly Leu Pro Ile
5 10 15

20 AGC ATG AAA ATT TTT ATG TAT TTA CTT ACT GTT TTT CTT ATC ACC CAG 150
Ser Met Lys Ile Phe Met Tyr Leu Leu Thr Val Phe Leu Ile Thr Gln
20 25 30 35

25 ATG ATT GGG TCA GCA CTT TTT GCT GTG TAT CTT CAT AGA AGG TTG GAC 198
Met Ile Gly Ser Ala Leu Phe Ala Val Tyr Leu His Arg Arg Leu Asp
40 45 50

30 AAG ATA GAA GAT GAA AGG AAT CTT CAT GAA GAT TTT GTA TTC ATG AAA 246
Lys Ile Glu Asp Glu Arg Asn Leu His Glu Asp Phe Val Phe Met Lys
55 60 65

35 ACG ATA CAG AGA TGC AAC ACA GGA GAA AGA TCC TTA TCC TTA CTG AAC 294
Thr Ile Gln Arg Cys Asn Thr Gly Glu Arg Ser Leu Ser Leu Leu Asn
70 75 80

40 TGT GAG GAG ATT AAA AGC CAG TTT GAA GGC TTT GTG AAG GAT ATA ATG 342
Cys Glu Glu Ile Lys Ser Gln Phe Glu Gly Phe Val Lys Asp Ile Met
85 90 95

45 TTA AAC AAA GAG GAG ACG AAG AAA GAA AAC AGC TTT GAA ATG CAA AAA 390
Leu Asn Lys Glu Glu Thr Lys Lys Glu Asn Ser Phe Glu Met Gln Lys
100 105 110 115

50 GGT GAT CAG AAT CCT CAA ATT GCG GCA CAT GTC ATA AGT GAG GCC AGC 438
Gly Asp Gln Asn Pro Gln Ile Ala Ala His Val Ile Ser Glu Ala Ser
120 125 130

55 AGT AAA ACA ACA TCT GTG TTA CAG TGG GCT GAA AAA GGA TAC TAC ACC 486
Ser Lys Thr Thr Ser Val Leu Gln Trp Ala Glu Lys Gly Tyr Tyr Thr
135 140 145

60 ATG AGC AAC AAC TTG GTA ACC CTG GAA AAT GGG AAA CAG CTG ACC GTT 534
Met Ser Asn Asn Leu Val Thr Leu Glu Asn Gly Lys Gln Leu Thr Val
150 155 160

AAA AGA CAA GGA CTC TAT TAT ATC TAT GCC CAA GTC ACC TTC TGT TCC 582
Lys Arg Gln Gly Leu Tyr Tyr Ile Tyr Ala Gln Val Thr Phe Cys Ser
165 170 175

AAT CGG GAA GCT TCG AGT CAA GCT CCA TTT ATA GCC AGC CTC TGC CTA 630
Asn Arg Glu Ala Ser Ser Gln Ala Pro Phe Ile Ala Ser Leu Cys Leu
180 185 190 195

AAG TCC CCC GGT AGA TTC GAG AGA ATC TTA CTC AGA GCT GCA AAT ACC 678
 Lys Ser Pro Gly Arg Phe Glu Arg Ile Leu Leu Arg Ala Ala Asn Thr
 200 205 210

5 CAC AGT TCC GCC AAA CCT TGC GGG CAA CAA TCC ATT CAC TTG GGA GGA 726
 His Ser Ser Ala Lys Pro Cys Gly Gln Gln Ser Ile His Leu Gly Gly
 215 220 225

10 GTA TTT GAA TTG CAA CCA GGT GCT TCG GTG TTT GTC AAT GTG ACT GAT 774
 Val Phe Glu Leu Gln Pro Gly Ala Ser Val Phe Val Asn Val Thr Asp
 230 235 240

CCA AGC CAA GTG AGC CAT GGC ACT GGC TTC ACG TCC TTT GGC TTA CTC 822
 Pro Ser Gln Val Ser His Gly Thr Gly Phe Thr Ser Phe Gly Leu Leu
 245 250 255

15 AAA CTC TGAACAGTGT CA 840
 Lys Leu
 260

20

(2) INFORMATION FOR SEQ ID NO:2:

- 25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 261 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear
 (ii) MOLECULE TYPE: protein
- 30 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Ile Glu Thr Tyr Asn Gln Thr Ser Pro Arg Ser Ala Ala Thr Gly
 1 5 10 15

35 Leu Pro Ile Ser Met Lys Ile Phe Met Tyr Leu Leu Thr Val Phe Leu
 20 25 30

Ile Thr Gln Met Ile Gly Ser Ala Leu Phe Ala Val Tyr Leu His Arg
 35 40 45

40 Arg Leu Asp Lys Ile Glu Asp Glu Arg Asn Leu His Glu Asp Phe Val
 50 55 60

45 Phe Met Lys Thr Ile Gln Arg Cys Asn Thr Gly Glu Arg Ser Leu Ser
 65 70 75 80

Leu Leu Asn Cys Glu Glu Ile Lys Ser Gln Phe Glu Gly Phe Val Lys
 85 90 95

50 Asp Ile Met Leu Asn Lys Glu Glu Thr Lys Lys Glu Asn Ser Phe Glu
 100 105 110

Met Gln Lys Gly Asp Gln Asn Pro Gln Ile Ala Ala His Val Ile Ser
 115 120 125

55 Glu Ala Ser Ser Lys Thr Thr Ser Val Leu Gln Trp Ala Glu Lys Gly
 130 135 140

60 Tyr Tyr Thr Met Ser Asn Asn Leu Val Thr Leu Glu Asn Gly Lys Gln
 145 150 155 160

Leu Thr Val Lys Arg Gln Gly Leu Tyr Tyr Ile Tyr Ala Gln Val Thr
 165 170 175
 5 Phe Cys Ser Asn Arg Glu Ala Ser Ser Gln Ala Pro Phe Ile Ala Ser
 180 185 190
 Leu Cys Leu Lys Ser Pro Gly Arg Phe Glu Arg Ile Leu Leu Arg Ala
 195 200 205
 10 Ala Asn Thr His Ser Ser Ala Lys Pro Cys Gly Gln Gln Ser Ile His
 210 215 220
 Leu Gly Gly Val Phe Glu Leu Gln Pro Gly Ala Ser Val Phe Val Asn
 225 230 235 240
 15 Val Thr Asp Pro Ser Gln Val Ser His Gly Thr Gly Phe Thr Ser Phe
 245 250 255
 20 Gly Leu Leu Lys Leu
 260

(2) INFORMATION FOR SEQ ID NO:3:

- 25 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 519 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 30 (ii) MOLECULE TYPE: cDNA
 (iii) HYPOTHETICAL: NO
 (iv) ANTI-SENSE: NO
 35 (vi) ORIGINAL SOURCE:
 (A) ORGANISM: HUMAN
 (vii) IMMEDIATE SOURCE:
 (B) CLONE: CD40 EXTRACELLULAR REGION
 40

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

45 CAGAACCACC CACTGCATGC AGAGAAAAAC AGTACCTAAT AAACAGTCAG TGCTGTTCTT 60
 TGTGCCAGCC AGGACAGAAA CTGGTGAGTG ACTGCACAGA GTTCACTGAA ACGGAATGCC 120
 TTCCTTGCGG TGAAAGCGAA TTCCTAGACA CCTGGAACAG AGAGACACAC TGCCACCAGC 180
 50 ACAAATACTG CGACCCCAAC CTAGGGCTTC GGGTCCAGCA GAAGGGCACC TCAGAAACAG 240
 ACACCATCTG CACCTGTGAA GAAGGCTGGC ACTGTACGAG TGAGGCCTGT GAGAGCTGTG 300
 TCCTGCACCG CTCATGCTCG CCCGGCTTTG GGGTCAAGCA GATTGCTACA GGGGTTTCTG 360
 55 ATACCATCTG CGAGCCCTGC CCAGTCGGCT TCTTCTCCAA TGTGTCATCT GCTTTCGAAA 420
 AATGTCACCC TTGGACAAGC TGTGAGACCA AAGACCTGGT TGTGCAACAG GCAGGCACAA 480
 60 ACAAGACTGA TGTGTCTGT GGTCCCCAGG ATCGGCTGA 519

(2) INFORMATION FOR SEQ ID NO:4:

5 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 740 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

10 (ii) MOLECULE TYPE: cDNA
 (iii) HYPOTHETICAL: NO
 (iv) ANTI-SENSE: NO

15 (vi) ORIGINAL SOURCE:
 (A) ORGANISM: HUMAN

(vii) IMMEDIATE SOURCE:
 (B) CLONE: IgG1 Fc

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

	CGGTACCGCT AGCGTCGACA GGCCTAGGAT ATCGATACGT AGAGCCCAGA TCTTGTGACA	60
25	AAACTCACAC ATGCCCACCG TGCCCAGCAC CTGAACTCCT GGGGGGACCG TCAGTCTTCC	120
	TCTTCCCCCC AAAACCCAAG GACACCCTCA TGATCTCCCG GACCCCTGAG GTCACATGCG	180
30	TGGTGGTGGG CGTGAGCCAC GAAGACCCTG AGGTCAAGTT CAACTGGTAC GTGGACGGCG	240
	TGGAGGTGCA TAATGCCAAG ACAAAGCCGC GGGAGGAGCA GTACAACAGC ACGTACCGGG	300
	TGGTCAGCGT CCTCACCGTC CTGCACCAGG ACTGGCTGAA TGGCAAGGAC TACAAGTGCA	360
35	AGGTCTCCAA CAAAGCCCTC CCAGCCCCCA TGCAGAAAAC CATCTCCAA GCCAAAGGGC	420
	AGCCCCGAGA ACCACAGGTG TACACCCTGC CCCCATCCCG GGATGAGCTG ACCAAGAACC	480
40	AGGTCAGCCT GACCTGCCTG GTCAAAGGCT TCTATCCCAG GCACATCGCC GTGGAGTGGG	540
	AGAGCAATGG GCAGCCGGAG AACAACTACA AGACCACGCC TCCCGTGCTG GACTCCGACG	600
	GCTCCTTCTT CCTCTACAGC AAGCTCACCG TGGACAAGAG CAGGTGGCAG CAGGGGAACG	660
45	TCTTCTCATG CTCCGTGATG CATGAGGCTC TGCACAACCA CTACACGCAG AAGAGCCTCT	720
	CCCTGTCTCC GGGTAAATGA	740

50 (2) INFORMATION FOR SEQ ID NO:5:

55 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 33 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Arg Met Lys Gln Ile Glu Asp Lys Ile Glu Glu Ile Leu Ser Lys Ile
 1 5 10 15
 Tyr His Ile Glu Asn Glu Ile Ala Arg Ile Lys Lys Leu Ile Gly Glu
 20 25 30
 Arg

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: not relevant
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Pro Asp Val Ala Ser Leu Arg Gln Gln Val Glu Ala Leu Gln Gly Gln
 1 5 10 15
 Val Gln His Leu Gln Ala Ala Phe Ser Gln Tyr
 20 25

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 929 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Human CD40L trimer

(ix) FEATURE:

- (A) NAME/KEY: sig_peptide
 (B) LOCATION: 65..142

(ix) FEATURE:

- (A) NAME/KEY: CDS
 (B) LOCATION: 65..886

(ix) FEATURE:

- (A) NAME/KEY: mat_peptide
 (B) LOCATION: 143..886

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

TGAGCGAGTC CGCATCGACG GATCGGAAAA CCTCTCCGAG GTACCTATCC CGGGGATCCC 60

	CACC	ATG	TTC	CAT	GTT	TCT	TTT	AGA	TAT	ATC	TTT	GGA	ATT	CCT	CCA	CTG	109
	Met	Phe	His	Val	Ser	Phe	Arg	Tyr	Ile	Phe	Gly	Ile	Pro	Pro	Leu		
	-26	-25					-20					-15					
5	ATC	CTT	GTT	CTG	CTG	CCT	GTC	ACT	AGT	TCT	GAC	CGT	ATG	AAA	CAG	ATA	157
	Ile	Leu	Val	Leu	Leu	Pro	Val	Thr	Ser	Ser	Asp	Arg	Met	Lys	Gln	Ile	
	-10						-5					1				5	
10	GAG	GAT	AAG	ATC	GAA	GAG	ATC	CTA	AGT	AAG	ATT	TAT	CAT	ATA	GAG	AAT	205
	Glu	Asp	Lys	Ile	Glu	Glu	Ile	Leu	Ser	Lys	Ile	Tyr	His	Ile	Glu	Asn	
					10					15					20		
15	GAA	ATC	GCC	CGT	ATC	AAA	AAG	CTG	ATT	GGC	GAG	CGG	ACT	AGT	TCT	GAC	253
	Glu	Ile	Ala	Arg	Ile	Lys	Lys	Leu	Ile	Gly	Glu	Arg	Thr	Ser	Ser	Asp	
				25					30					35			
20	AAG	ATA	GAA	GAT	GAA	AGG	AAT	CTT	CAT	GAA	GAT	TTT	GTA	TTC	ATG	AAA	301
	Lys	Ile	Glu	Asp	Glu	Arg	Asn	Leu	His	Glu	Asp	Phe	Val	Phe	Met	Lys	
			40					45					50				
25	ACG	ATA	CAG	AGA	TGC	AAC	ACA	GGA	GAA	AGA	TCC	TTA	TCC	TTA	CTG	AAC	349
	Thr	Ile	Gln	Arg	Cys	Asn	Thr	Gly	Glu	Arg	Ser	Leu	Ser	Leu	Leu	Asn	
		55					60					65					
30	TGT	GAG	GAG	ATT	AAA	AGC	CAG	TTT	GAA	GGC	TTT	GTG	AAG	GAT	ATA	ATG	397
	Cys	Glu	Glu	Ile	Lys	Ser	Gln	Phe	Glu	Gly	Phe	Val	Lys	Asp	Ile	Met	
		70				75					80					85	
35	TTA	AAC	AAA	GAG	GAG	ACG	AAG	AAA	GAA	AAC	AGC	TTT	GAA	ATG	CAA	AAA	445
	Leu	Asn	Lys	Glu	Glu	Thr	Lys	Lys	Glu	Asn	Ser	Phe	Glu	Met	Gln	Lys	
					90					95					100		
40	GGT	GAT	CAG	AAT	CCT	CAA	ATT	GCG	GCA	CAT	GTC	ATA	AGT	GAG	GCC	AGC	493
	Gly	Asp	Gln	Asn	Pro	Gln	Ile	Ala	Ala	His	Val	Ile	Ser	Glu	Ala	Ser	
				105					110					115			
45	AGT	AAA	ACA	ACA	TCT	GTG	TTA	CAG	TGG	GCT	GAA	AAA	GGA	TAC	TAC	ACC	541
	Ser	Lys	Thr	Thr	Ser	Val	Leu	Gln	Trp	Ala	Glu	Lys	Gly	Tyr	Tyr	Thr	
			120					125					130				
50	ATG	AGC	AAC	AAC	TTG	GTA	ACC	CTG	GAA	AAT	GGG	AAA	CAG	CTG	ACC	GTT	589
	Met	Ser	Asn	Asn	Leu	Val	Thr	Leu	Glu	Asn	Gly	Lys	Gln	Leu	Thr	Val	
		135					140					145					
55	AAA	AGA	CAA	GGA	CTC	TAT	TAT	ATC	TAT	GCC	CAA	GTC	ACC	TTC	TGT	TCC	637
	Lys	Arg	Gln	Gly	Leu	Tyr	Tyr	Ile	Tyr	Ala	Gln	Val	Thr	Phe	Cys	Ser	
		150					155				160					165	
60	AAT	CGG	GAA	GCT	TCG	AGT	CAA	GCT	CCA	TTT	ATA	GCC	AGC	CTC	TGC	CTA	685
	Asn	Arg	Glu	Ala	Ser	Ser	Gln	Ala	Pro	Phe	Ile	Ala	Ser	Leu	Cys	Leu	
					170					175					180		
65	AAG	TCC	CCC	GGT	AGA	TTC	GAG	AGA	ATC	TTA	CTC	AGA	GCT	GCA	AAT	ACC	733
	Lys	Ser	Pro	Gly	Arg	Phe	Glu	Arg	Ile	Leu	Leu	Arg	Ala	Ala	Asn	Thr	
			185						190					195			
70	CAC	AGT	TCC	GCC	AAA	CCT	TGC	GGG	CAA	CAA	TCC	ATT	CAC	TTG	GGA	GGA	781
	His	Ser	Ser	Ala	Lys	Pro	Cys	Gly	Gln	Gln	Ser	Ile	His	Leu	Gly	Gly	
			200					205					210				

	GTA TTT GAA TTG CAA CCA GGT GCT TCG GTG TTT GTC AAT GTG ACT GAT	829
	Val Phe Glu Leu Gln Pro Gly Ala Ser Val Phe Val Asn Val Thr Asp	
	215 220 225	
5	CCA AGC CAA GTG AGC CAT GGC ACT GGC TTC ACG TCC TTT GGC TTA CTC	877
	Pro Ser Gln Val Ser His Gly Thr Gly Phe Thr Ser Phe Gly Leu Leu	
	230 235 240 245	
10	AAA CTC TGAGCGGCCG CTACAGATGA ATAATAAGCA TGTTCGGATT CCTCAA	929
	Lys Leu	

(2) INFORMATION FOR SEQ ID NO:8:

15

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 273 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

25	Met Phe His Val Ser Phe Arg Tyr Ile Phe Gly Ile Pro Pro Leu Ile	
	-26 -25 -20 -15	
	Leu Val Leu Leu Pro Val Thr Ser Ser Asp Arg Met Lys Gln Ile Glu	
	-10 -5 1 5	
30	Asp Lys Ile Glu Glu Ile Leu Ser Lys Ile Tyr His Ile Glu Asn Glu	
	10 15 20	
35	Ile Ala Arg Ile Lys Lys Leu Ile Gly Glu Arg Thr Ser Ser Asp Lys	
	25 30 35	
	Ile Glu Asp Glu Arg Asn Leu His Glu Asp Phe Val Phe Met Lys Thr	
	40 45 50	
40	Ile Gln Arg Cys Asn Thr Gly Glu Arg Ser Leu Ser Leu Leu Asn Cys	
	55 60 65 70	
	Glu Glu Ile Lys Ser Gln Phe Glu Gly Phe Val Lys Asp Ile Met Leu	
	75 80 85	
45	Asn Lys Glu Glu Thr Lys Lys Glu Asn Ser Phe Glu Met Gln Lys Gly	
	90 95 100	
50	Asp Gln Asn Pro Gln Ile Ala Ala His Val Ile Ser Glu Ala Ser Ser	
	105 110 115	
	Lys Thr Thr Ser Val Leu Gln Trp Ala Glu Lys Gly Tyr Tyr Thr Met	
	120 125 130	
55	Ser Asn Asn Leu Val Thr Leu Glu Asn Gly Lys Gln Leu Thr Val Lys	
	135 140 145 150	
	Arg Gln Gly Leu Tyr Tyr Ile Tyr Ala Gln Val Thr Phe Cys Ser Asn	
	155 160 165	
60		

	Arg	Glu	Ala	Ser	Ser	Gln	Ala	Pro	Phe	Ile	Ala	Ser	Leu	Cys	Leu	Lys
				170					175					180		
5	Ser	Pro	Gly	Arg	Phe	Glu	Arg	Ile	Leu	Leu	Arg	Ala	Ala	Asn	Thr	His
			185					190					195			
	Ser	Ser	Ala	Lys	Pro	Cys	Gly	Gln	Gln	Ser	Ile	His	Leu	Gly	Gly	Val
		200					205					210				
10	Phe	Glu	Leu	Gln	Pro	Gly	Ala	Ser	Val	Phe	Val	Asn	Val	Thr	Asp	Pro
	215					220					225					230
	Ser	Gln	Val	Ser	His	Gly	Thr	Gly	Phe	Thr	Ser	Phe	Gly	Leu	Leu	Lys
					235					240					245	
15	Leu															

CLAIMS

What is claimed is:

1. An isolated DNA encoding a CD40L mutein, selected from the group consisting
5 of:
 (a) a DNA that encodes a polypeptide having an amino acid sequence as set forth in
 SEQ ID NO:2 wherein a cysteine at amino acid 194 is replaced with another
 amino acid; and
 (b) a DNA that encodes a polypeptide that is a fragment of the mutein (a) and which
10 polypeptide binds CD40;
wherein the amino acid that is substituted for the cysteine at amino acid 194 is selected from
the group consisting of tryptophan, serine, aspartic acid, and lysine.

2. The isolated DNA of claim 1 that encodes a polypeptide selected from the group
15 consisting of polypeptides comprising amino acids 1 through 261, 35 through 261, 34
through 225, 113 through 261, 113 through 225, 120 through 261, or 120 through 225 of
SEQ ID NO:2.

3. The isolated DNA according to claim 2 which further comprises a DNA
20 encoding an oligomerizing zipper having an amino acid sequence represented by SEQ ID
NO:5.

4. The isolated DNA of claim 3, wherein the DNA encodes a fusion protein
comprising the oligomerizing zipper fused to the amino-proximal end of a fragment of the
25 extracellular domain of CD40L consisting of amino acids 113 through 261 of SEQ ID
NO:2.

5. A recombinant expression vector comprising a DNA according to claim 4.

- 30 6. A host cell transformed or transfected with a recombinant expression vector
according to claim 5.

7. A process for preparing a CD40L polypeptide, comprising culturing a host cell
according to claim 6 under conditions promoting expression and recovering CD40L
35 polypeptide from the culture.

8. A CD40L polypeptide encoded by a DNA according to any one of claims 1
through 4.

1/2

Binding of human and Murine CD40 LT and CD40 L Fc
Dimer to CD40Fc by Biosensor Assay





FIG. 1

2/2

Binding of CD40 ligands to CD40Fc using equilibrium binding values estimated from a kinetic analysis of the association phase

**FIG. 2A****FIG. 2B**

This Page Blank (uspto)

INTERNATIONAL SEARCH REPORT

International Application No

PC1/US 96/09632

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C07K14/725 C12N15/62 C12N5/10 C12N1/19

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO,A,93 08207 (IMMUNEX CORPORATION) 29 April 1993 cited in the application see the whole document ---	1,2
A	WO,A,94 10308 (IMMUNEX CORPORATION) 11 May 1994 see example 1 see page 4, line 20 - page 6, line 17 --- -/--	1-8

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

G document member of the same patent family

Date of the actual completion of the international search

26 November 1996

Date of mailing of the international search report

11 12 96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-2016

Authorized officer

Le Cornec. N

INTERNATIONAL SEARCH REPORT

International Application No

PCI/US 96/09632

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	BIOCHEMISTRY, vol. 34, no. 6, February 1995, EASTON, PA US, pages 1833-1844, XP002019526 J. BAJORATH ET AL: "Identification of residues on CD40 and its ligand which are critical for the receptor-ligand interaction" see the whole document ---	1
A	EP,A,0 585 943 (BRISTOL-MYERS SQUIBB COMPANY) 9 March 1994 see page 6, line 26 - page 8, line 27 see page 11, line 1 - line 48 -----	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 96/09632

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9308207	29-04-93	AU-B- 661360	20-07-95
		AU-A- 3122693	21-05-93
		CA-A- 2121798	29-04-93
		EP-A- 0667901	23-08-95
		FI-A- 941837	30-05-94
		JP-T- 7504083	11-05-95
		NO-A- 941422	27-06-94

WO-A-9410308	11-05-94	AU-A- 5537794	24-05-94
		CA-A- 2146559	11-05-94
		EP-A- 0672141	20-09-95
		JP-T- 8507676	20-08-96
		NZ-A- 257942	26-04-96

EP-A-585943	09-03-94	US-A- 5540926	30-07-96
		AU-A- 4612093	10-03-94
		CA-A- 2105552	05-03-94
		FI-A- 933862	05-03-94
		HU-A- 69977	28-09-95
		JP-A- 6315383	15-11-94
		NO-A- 933126	07-03-94
		NZ-A- 248569	26-10-95
ZA-A- 9306491	25-03-94		

This Page Blank (uspto)